

# The Effect of Sequential Contrast Agent Studies on the Assessment of Relative Cerebral Blood Volume

A. P. Pathak<sup>1,2</sup>, P. A. Bandettini<sup>2,1</sup>, R. Risinger<sup>3</sup>, E. Stein<sup>3</sup>, and K. M. Donahue<sup>2,1</sup>

<sup>1</sup>Department of Biomedical Engineering, Marquette University, Milwaukee, Wisconsin, U. S. A.

<sup>2</sup>Biophysics Research Institute, <sup>3</sup>Dept. of Psychiatry, Medical College of Wisconsin, Milwaukee, Wisconsin, U. S. A.

**INTRODUCTION.** Dynamic MR imaging of the first-pass of a susceptibility contrast agent confined to the intravascular space has enabled the creation of relative cerebral blood volume (rCBV) maps. This approach has been used for the evaluation of brain neoplasms [1] and for the detection of brain activation changes [2]. Comparison of rCBV for cortical gray versus periventricular white matter yielded a ratio of two, a value that has been well documented in the literature, thus supporting the accuracy of this approach [3]. Recently though, the stability and reproducibility of this method for sequential rCBV measurements has been called into question with results indicating artifactually elevated estimates of rCBV with subsequent contrast agent doses [4]. This has important implications for the utility of susceptibility-rCBV methods to evaluate changes in blood volume that might occur between resting and activated states. However, the results described in [4], were derived using a whole brain region of interest (ROI). Whether the ratio of gray to white matter rCBV remained constant with subsequent contrast agent doses was not addressed, and was the primary goal of the current study.

**METHODS.** To study the effect of multiple doses of contrast agent, two normal subjects were studied with three sequential boluses of Gd. All images were obtained using a Bruker BIOSPEC 3T imaging system. Interleaved gradient echo (64x64, TR=1s, TE=27.2 ms) and spin echo images (64x64, TR=1s, TE=110 ms) were acquired in the axial plane at, 7 mm thick slices, over 180 seconds. Sixty seconds into the acquisition, 0.05 mmol/kg Gd-DTPA, i.e. Prohance<sup>®</sup> was bolus (approx. 6-8s) injected through a 12-gauge venous catheter. From 1-12 mins later, a second injection was given, followed by a third after the same time interval, both 0.05 mmol/kg.

The rCBV for each pixel was computed according to the following relationship:

$$rCBV \propto \frac{-1}{TE} \int \ln \frac{S(t)}{S(0)} dt \quad (1)$$

where  $S(t)$  is the MR signal intensity,  $S(0)$  is the averaged baseline signal intensity, and  $TE$  is the echo-time. The initial point for the integration was determined on the basis of a gamma-variate fit to the concentration-time image data. The gamma-variate function with a recirculation cut-off is defined as follows:

$$S(t) = Q(t - t_0)^r e^{-(t-t_0)^b} \quad (2)$$

where  $S(t)$  is the MR signal intensity,  $Q$ ,  $r$ ,  $b$  are fit constants,  $t$  is the injection time, and  $t_0$  is the appearance time of the tracer. Integration limits were set from  $t_0$  to  $(t_0+12)$  points. For the whole brain CBV a ROI of 25x25 voxels was used and histogram analysis was carried, to assess the CBV distribution on a voxel-wise basis. Finally, for the gray/white ratio study, ROIs were selected in cortical gray and periventricular white matter from high-resolution anatomical images.

**RESULTS.** For the range of doses studied, we observed changes in the whole brain CBV as a function of contrast agent dose (Table 1.), similar to previous studies using whole brain [4]. However, a histogram analysis of the gradient and spin echo rCBVs (Fig. 1, Fig. 2) revealed that the relative distribution of blood-volume as a function of contrast agent dose, remains unchanged on a voxel-wise basis, for both gradient and spin echo sequences. Finally, single factor ANOVA reveals that the

gray/white matter ratios are not significantly different from dose to dose (Fig. 3).

Table 1.

Sequence	Dose	Subject 1		Subject 2	
		rCBV	ΔrCBV	rCBV	ΔrCBV
GE	1	7399	-	4628	-
	2	6200	-16.2%	11559	-150.6%
	3	6752	8.90%	11935	3.25%
SE	1	5729	-	1265	-
	2	4237	-26.0%	2740	116.6%
	3	5814	37.2%	4323	57.8%




Figure 1.




Figure 2.




Figure 3.

**DISCUSSION/CONCLUSION.** This study demonstrates that the utility of sequential contrast agent studies is not limited to computing the rCBV only after attaining a relative steady state after multiple boluses [4]. We have demonstrated that the resting blood volume ratio of gray to white matter stays constant from bolus to bolus, enabling us to evaluate changes in gray matter rCBV when normalized to white matter rCBV. Such might occur between resting and activated states during brain activation studies. However, this is still not an appropriate method for detecting rCBV changes when both, gray and white matter rCBV are increased, as in the case of hypercapnia studies.

## References

1. Aronen, H. J. et al., *Radiology* 191:41-51, 1994.
2. Belliveau, J. W. et al., *Science* 254:716-719, 1991.
3. Belliveau, J. W. et al., *Magn. Reson. Med.* 14:538-546, 1990.
4. Levin, J. M. et al., *Magn. Reson. Med.* 34:655-663, 1995.

1998 Abstract Form for Scientific Presentations  
INTERNATIONAL SOCIETY FOR  
MAGNETIC RESONANCE IN MEDICINE  
SIXTH SCIENTIFIC MEETING  
SYDNEY AUSTRALIA  
APRIL 18-24, 1998

## ABSTRACT DEADLINE:

Received no later than NOVEMBER 18, 1997.  
All copyrights to accepted abstracts become the property of the ISMRM; no proprietary information may be withheld by authors.

If the authors have submitted their abstract electronically an additional fee must be paid.

☐ Prefer Oral Presentation but willing to present as a poster

☐ Video required (available only for Oral Presentations)

☐ Prefer Poster but willing to make Oral Presentation

☐ Poster only

☐ The authors certify that any work with human or animal subjects complies with the guiding policies of their national or institutional regulatory bodies.

Topic categories (please fill in both):

Enter "old" category number:

Enter "new" category numbers: A  B  C  D  E

Keywords: rCBV susceptibility contrast sequential

SEND BACK TO FORM FOR ALL INSTRUCTIONS

Please type the name and complete mailing address of the first author

Name: Arvind Pathak

Address: Biophysics Res. Inst., Medical College of WI, 8701 Watertown Plank Rd., Milwaukee, WI 53226

Country: U.S.A.

Telephone: (414) 456-4051

FAX: (414) 456-4512

e-mail: pathakap@post.its.mcw.edu